

Tapering plan for desvenlafaxine – ‘Even slower’ taper

This plan gives step-by-step instructions to slowly decrease and stop antidepressants. Following a plan like this can help to minimise antidepressant withdrawal symptoms. **Take this tapering plan to discuss with your doctor.**

- Aim to reduce your antidepressant dose every **2-4 weeks**. The tapering speed is **flexible** – you can pause, go more slowly or more quickly depending on any withdrawal symptoms you may experience.
- If you experience withdrawal symptoms, you can return to your previous dose for symptom relief. When ready to try again, you could decrease your dose in even smaller reductions and leave longer between dose reductions.
- Do NOT skip the final steps with small dose reductions – these are very important to prevent withdrawal symptoms.
- Do not skip a day or alternate days or doses. Do not suddenly stop antidepressants.
- Desvenlafaxine has higher risk for withdrawal symptoms. Read about withdrawal symptoms on the next page.

Availability: Desvenlafaxine (Pristiq, Desfax, Desven) is available in 50mg and 100mg extended-release tablets. Mini doses are available in capsules on prescription from a compounding chemist. The manufacturer advises against dividing, crushing, chewing, or dissolving desvenlafaxine tablets. eMIMS: <https://app-emimselite-com.ap1.proxy.openathens.net/Drtc?id=116>

Alternatively, switch to venlafaxine for tapering, as the pharmacology is similar. Desvenlafaxine 100mg = venlafaxine 150mg.

Step	Daily dose	Daily capsules	Date I started this dose	Tick when done	Step	Daily dose	Daily capsules	Date I started this dose	Tick when done
1	200mg	2 x 100mg tablets			21	15mg	1 x 10mg & 1 x 5mg		
2	150mg	3 x 50mg tablets			22	13mg	1 x 10mg & 3 x 1mg		
3	100mg	2 x 50mg tablets			23	12mg	1 x 10mg & 1 x 2mg		
4	90mg	1 x 50mg tablet & 4 x 10mg capsules			24	10mg	1 x 10mg		
5	80mg	1 x 50mg tablet & 3 x 10mg capsules			25	9mg	1 x 5mg & 2 x 2mg		
6	70mg	1 x 50mg tablet & 2 x 10mg capsules			26	8mg	1 x 5mg & 3 x 1mg		
7	60mg	1 x 50mg tablet & 1 x 10mg capsule			27	7mg	1 x 5mg & 1 x 2mg		
8	55mg	1 x 50mg tablet & 1 x 5mg capsule			28	6mg	1 x 5mg & 1 x 1mg		
9	50mg	1 x 50mg tablet			29	5mg	1 x 5mg		
10	45mg	2 x 20mg & 1 x 5mg			30	4.5mg	2 x 2mg & 1 x 0.5mg		
11	40mg	2 x 20mg			31	4mg	2 x 2mg		
12	35mg	3 x 10mg & 1 x 5mg			32	3.5mg	1 x 2mg & 3 x 0.5mg		
13	30mg	3 x 10mg			33	3mg	1 x 2mg & 1 x 1mg		
14	27mg	1 x 20mg & 1 x 5mg & 1 x 2mg			34	2.5mg	1 x 2mg & 1 x 0.5mg		
15	25mg	1 x 20mg & 1 x 5mg			35	2mg	1 x 2mg		
16	22mg	1 x 20mg & 1 x 2mg			36	1.5mg	1 x 1mg & 1 x 0.5mg		
17	20mg	1 x 20mg			37	1mg	1 x 1mg		
18	19mg	3 x 5mg & 2 x 2mg			38	0.5mg	1 x 0.5mg		
19	17mg	3 x 5mg & 1 x 2mg			39	STOP			
20	16mg	3 x 5mg & 1 x 1mg							

These steps are most important. Do not skip.

Is it withdrawal or is it relapse?

It can be challenging weaning off antidepressants. You may be feeling anxious, irritable and tearful and wondering whether you are experiencing withdrawal or relapse. It is not always easy to tell.

Antidepressant withdrawal symptoms are common, and more likely if you have been taking an antidepressant for a long time or at higher doses.

Antidepressant withdrawal symptoms

Emotional	anxiety/nervousness, worsened mood, irritability, bouts of crying, mood swings, anger outbursts, feelings of panic
Physical	dizziness/light-headedness, fatigue/tiredness, sensation of spinning/vertigo, insomnia, agitation, feeling of inner restlessness, brain 'zaps' or electric shock sensations, headache, rapid heartbeat, nausea, diarrhoea, muscle cramps, unsteady gait, tinnitus
Cognitive	brain fog, increased sensitivity to light or sound, memory problems, vivid dreams

Distinguishing withdrawal from relapse

Maintain a high index of suspicion. A good rule of thumb is that if new symptoms appear when you are decreasing your dose then 'the symptoms are likely to be withdrawal until proven otherwise'.

If you are experiencing physical symptoms, such as dizziness or light-headedness, that were never part of your original problem then you are likely experiencing withdrawal symptoms.

If your symptoms are relieved on reverting to a higher dose, then it is likely to be withdrawal.

Generally, relapse (depression or anxiety) takes longer to manifest and to respond to treatment.

For supporting evidence and further information see www.releasetoolkit.com.au or [click here](#) or scan below



How long do withdrawal symptoms last?

Withdrawal symptoms often gradually subside over weeks, reflecting the time that it takes for the brain and body to adapt to being without antidepressants (not the antidepressant half-life). Unfortunately, for some people, withdrawal symptoms can be severe and protracted.

How to relieve withdrawal symptoms

Returning to your previous dose will often relieve symptoms. When ready to try again, decrease your dose even more slowly, making smaller reductions and leaving longer between reductions, for example decreasing by 5% every few weeks or months.

How to minimise withdrawal symptoms

The best way to minimise withdrawal symptoms is by slowly decreasing your dose, through to very low doses. This can take many months. You may think that the low doses in hyperbolic tapering are not needed, but small dose changes can have large effects. Do NOT skip the second half of the tapering plan. Do NOT skip a day or alternate days or doses, and do not suddenly stop.

Build a support team around you, as you may not recognise withdrawal symptoms in yourself. Look after yourself. Try physical activity and relaxation, sleep and eat well, limit alcohol and other drugs.

Why stop antidepressants?

For the health and quality of life benefits. You may feel more alert and energetic and you may begin to feel your emotions again. You may find it easier to manage your weight, and your sexual functioning may improve. You may prefer living your life free of antidepressant medication.

There are benefits in reducing your dose, even if you cannot stop. It is not a failure if you do not manage to stop. You can always try again at some point in the future, tapering even more slowly.

