

Tapering plan for duloxetine – ‘Even slower’ taper

This plan gives step-by-step instructions to slowly decrease and stop antidepressants. Following a plan like this can help to minimise antidepressant withdrawal symptoms. **Take this tapering plan to discuss with your doctor.**

- Aim to reduce your antidepressant dose every **2-4 weeks**. The tapering speed is **flexible** – you can pause, go more slowly or more quickly depending on any withdrawal symptoms you may experience.
- If you experience withdrawal symptoms, you can return to your previous dose for symptom relief. When ready to try again, you could decrease your dose in even smaller reductions and leave longer between dose reductions.
- Do NOT skip the final steps with small dose reductions – these are very important to prevent withdrawal symptoms.
- Do not skip a day or alternate days or doses. Do not suddenly stop antidepressants.
- Duloxetine has higher risk for withdrawal symptoms. Read about withdrawal symptoms on the next page.

Availability: Duloxetine (Cymbalta, Andepra, Duloxecor, Dytrex, Tixel) is available in 30mg and 60mg capsules containing enteric coated beads. Mini dose capsules are available on prescription from a compounding pharmacy OR by opening a capsule, counting or weighing beads using clean dry hands and tweezers or a ruler, and putting dose of beads back into the capsule or another gelatine capsule bought from a pharmacy or online for ingestion. The manufacturer advises that capsules may be opened and beads stored in a jar as their enteric coating is stable in air. The beads should not be swallowed without a capsule and should not be chewed or dispersed in water. eMIMS: <https://app-emimselite.com.ap1.proxy.openathens.net/Drct?id=140>

Beads: Each capsule contains the same weight of drug but because the beads vary in size, capsules may contain different numbers of beads. If your 30mg capsule contains around 250 beads, then a 5mg dose is around 42 beads.

Step	Daily dose	Daily capsules	Date I started this dose	Tick when done	Step	Daily dose	Daily capsules	Date I started this dose	Tick when done
1	60mg	2 x 30mg capsules			21	5.2mg	1 x 5mg & 1 x 0.2mg capsules		
2	50mg	1 x 30mg & 2 x 10mg capsules			22	4.6mg	2 x 2mg & 2 x 0.3mg capsules		
3	40mg	1 x 30mg & 1 x 10mg capsules			23	4.3mg	2 x 2mg & 1 x 0.3mg capsules		
4	35mg	1 x 30mg & 1 x 5mg capsules			24	3.9mg	1 x 3mg & 3 x 0.3mg capsules		
5	30mg	1 x 30mg capsule			25	3.5mg	1 x 3mg & 1 x 0.5mg capsules		
6	25mg	2 x 10mg & 1 x 5mg capsules			26	3.15mg	1 x 3mg & 1 x 0.15mg capsules		
7	22mg	2 x 10mg & 1 x 2mg capsules			27	2.9mg	1 x 2mg & 3 x 0.3mg capsules		
8	20mg	2 x 10mg capsules			28	2.5mg	2 x 1mg & 1 x 0.5mg capsules		
9	18mg	1 x 10mg & 4 x 2mg capsules			29	2.2mg	1 x 2mg & 1 x 0.2mg capsules		
10	16mg	1 x 10mg & 3 x 2mg capsules			30	2mg	2 x 1mg capsules		
11	14mg	1 x 10mg & 2 x 2mg capsules			31	1.7mg	3 x 0.5mg & 1 x 0.2mg capsules		
12	13mg	1 x 10mg & 3 x 1mg capsules			32	1.5mg	3 x 0.5mg capsules		
13	11mg	1 x 10mg & 1 x 1mg capsules			33	1.3mg	1 x 1mg & 2 x 0.15mg capsules		
14	10mg	1 x 10mg capsules			34	1mg	1 x 1mg capsule		
15	9mg	3 x 3mg capsules			35	0.8mg	4 x 0.2mg capsules		
16	8.3mg	1x 5mg, 1x 3mg, 1x 0.3mg capsules			36	0.65mg	1x 0.5mg & 1 x 0.15mg capsules		
17	7.6mg	1x 5mg, 1x 2mg, 2 x 0.3mg capsules			37	0.5mg	1 x 0.5mg capsule		
18	7mg	1 x 5mg & 1 x 2mg capsules			38	0.3mg	2 x 0.15mg capsules		
19	6.3mg	2 x 3mg & 1 x 0.3mg capsules			39	0.15mg	1 x 0.15mg capsule		
20	5.6mg	1 x 5mg & 2 x 0.3mg capsules			40	STOP			

These steps are most important. Do not skip.

Is it withdrawal or is it relapse?

It can be challenging weaning off antidepressants. You may be feeling anxious, irritable and tearful and wondering whether you are experiencing withdrawal or relapse. It is not always easy to tell.

Antidepressant withdrawal symptoms are common, and more likely if you have been taking an antidepressant for a long time or at higher doses.

Antidepressant withdrawal symptoms

Emotional	anxiety/nervousness, worsened mood, irritability, bouts of crying, mood swings, anger outbursts, feelings of panic
Physical	dizziness/light-headedness, fatigue/tiredness, sensation of spinning/vertigo, insomnia, agitation, feeling of inner restlessness, brain 'zaps' or electric shock sensations, headache, rapid heartbeat, nausea, diarrhoea, muscle cramps, unsteady gait, tinnitus
Cognitive	brain fog, increased sensitivity to light or sound, memory problems, vivid dreams

Distinguishing withdrawal from relapse

Maintain a high index of suspicion. A good rule of thumb is that if new symptoms appear when you are decreasing your dose then 'the symptoms are likely to be withdrawal until proven otherwise'.

If you are experiencing physical symptoms, such as dizziness or light-headedness, that were never part of your original problem then you are likely experiencing withdrawal symptoms.

If your symptoms are relieved on reverting to a higher dose, then it is likely to be withdrawal.

Generally, relapse (depression or anxiety) takes longer to manifest and to respond to treatment.

For supporting evidence and further information see www.releasetoolkit.com.au or [click here](#) or scan below



How long do withdrawal symptoms last?

Withdrawal symptoms often gradually subside over weeks, reflecting the time that it takes for the brain and body to adapt to being without antidepressants (not the antidepressant half-life). Unfortunately, for some people, withdrawal symptoms can be severe and protracted.

How to relieve withdrawal symptoms

Returning to your previous dose will often relieve symptoms. When ready to try again, decrease your dose even more slowly, making smaller reductions and leaving longer between reductions, for example decreasing by 5% every few weeks or months.

How to minimise withdrawal symptoms

The best way to minimise withdrawal symptoms is by slowly decreasing your dose, through to very low doses. This can take many months. You may think that the low doses in hyperbolic tapering are not needed, but small dose changes can have large effects. Do NOT skip the second half of the tapering plan. Do NOT skip a day or alternate days or doses, and do not suddenly stop.

Build a support team around you, as you may not recognise withdrawal symptoms in yourself. Look after yourself. Try physical activity and relaxation, sleep and eat well, limit alcohol and other drugs.

Why stop antidepressants?

For the health and quality of life benefits. You may feel more alert and energetic and you may begin to feel your emotions again. You may find it easier to manage your weight, and your sexual functioning may improve. You may prefer living your life free of antidepressant medication.

There are benefits in reducing your dose, even if you cannot stop. It is not a failure if you do not manage to stop. You can always try again at some point in the future, tapering even more slowly.

