RELEASE
Redressing long-term antidepressant use

The University Of Queensland
AUSTRALIA
CREATE CHANGE

Tapering plan for vortioxetine – 'Faster' taper

This plan gives step-by-step instructions to slowly decrease and stop antidepressants. Following a plan like this can help to minimise antidepressant withdrawal symptoms.

Take this tapering plan to discuss with your doctor.

- Aim to reduce your antidepressant dose every **2-4 weeks**. The tapering speed is **flexible** – you can pause, go more slowly or more quickly depending on any withdrawal symptoms you may experience.
- If you experience withdrawal symptoms, you can return to your previous dose for symptom relief. When ready to try again, you could decrease your dose in even smaller reductions and leave longer between dose reductions.
- Do NOT skip the final steps with small dose reductions – these are very important to prevent withdrawal symptoms.
- Do not skip a day or alternate days or doses. Do not suddenly stop antidepressants.
- Vortioxetine has moderate risk for withdrawal symptoms. Read about withdrawal symptoms on the next page.

Vortioxetine (Trintellix and Brintellix) is available in 5mg, 10mg, 15mg, and 20mg tablet. Mini doses are available in capsules on prescription from a compounding chemist or by making your own liquid suspension following the instructions below.

Step	Daily dose		Taper using Liquid or Capsules		Date (I started this step)	Tick when completed
			Liquid option (1mg/ml)	Capsules option		
1.		**20mg**	4 x 5mg tablets			
2.		**15mg**	3 x 5mg tablets			
3.		**10mg**	2 x 5mg tablets			
4.		**7.5mg**	1½ x 5mg tablets			
5.		**5mg**	1 x 5mg tablet			
6.	These steps are most important. Do not skip.	**3.5mg**	3.5ml liquid	½ x 5mg tablet & 2 x 0.5mg capsules		
7.		**2.5mg**	½ x 5mg tablet			
8.		**1.5mg**	1.5ml liquid	3 x 0.5mg capsules		
9.		**1mg**	1ml liquid	2 x 0.5mg capsules		
10.		**0.5mg**	0.5ml liquid	1 x 0.5mg capsule		
11.	STOP					

Acknowledgement: Dr Mark Horowitz for guidance on dosing schedule (see Maudsley Deprescribing Guidelines for more information). 4 Aug 2026

How to make vortioxetine liquid suspension: You will get 5ml of 1mg/ml liquid suspension

1. You will need a pill cutter, 5ml and 1ml syringes, and a jar (or similar) for mixing.
2. Crush a 5mg tablet between two teaspoons over a small plate.
3. Place crushed tablet into 5ml water in a jar. Shake or stir vigorously to spread the drug evenly through the mixture. The liquid may be cloudy because some parts of the tablet (the outer covering) do not dissolve.
4. Measure prescribed daily liquid using a syringe and consume.
5. Make this liquid suspension fresh daily. Discard any unused into the rubbish.

Pharmaceutical guidance advises that vortioxetine tablets can be crushed and dispersed in water for administration eMIMS: https://app-emimselite-com.ap1.proxy.openathens.net/Drtc?id=1064 (UK NEWT guidelines, NHS pharmaceutical guidance). www.sps.nhs.uk/articles/ssri-suggestions-for-adults-with-swallowing-difficulties/

Video demonstrating how to crush and disperse tablets, and measure liquid: https://vimeo.com/1170454505?share=copy&fl=sv&fe=ci

RELEASE
Redressing long-term antidepressant use

The University Of Queensland
AUSTRALIA
CREATE CHANGE

Is it withdrawal or relapse?

The body adapts to being on antidepressants and, when stopping, must re-adapt. This takes time and is why hyperbolic tapering may be necessary.

Antidepressant withdrawal symptoms are common, particularly if you have been taking antidepressants for a long time or at higher doses. You may be feeling anxious, irritable and tearful and wondering whether you are experiencing withdrawal symptoms or relapse.

Antidepressant withdrawal symptoms

Emotional	anxiety/nervousness, worsened mood, irritability, bouts of crying, mood swings, anger outbursts, feelings of panic
Physical	dizziness/light-headedness, fatigue/tiredness, sensation of spinning/vertigo, insomnia, agitation, feeling of inner restlessness, brain 'zaps' or electric shock sensations, headache, rapid heartbeat, nausea, diarrhoea, muscle cramps, unsteady gait, tinnitus
Cognitive	brain fog, increased sensitivity to light or sound, memory problems, vivid dreams

Distinguishing withdrawal from relapse

Maintain a high index of suspicion. A good rule of thumb is that if new symptoms appear when you are decreasing your dose then 'the symptoms are likely to be withdrawal until proven otherwise'.

If you are experiencing physical symptoms, such as dizziness or light-headedness, that were never part of your original problem then you are likely experiencing withdrawal symptoms.

If your symptoms are relieved on reverting to a higher dose, then it is likely to be withdrawal.

Generally, relapse (depression or anxiety) takes longer to manifest and to respond to treatment.

For supporting evidence and further information see www.releasetoolkit.com.au or click here or scan below

How long do withdrawal symptoms last?

Withdrawal symptoms often gradually subside over weeks, reflecting the time that it takes for the brain and body to adapt to being without antidepressants (not the antidepressant half-life). Unfortunately, for some people, withdrawal symptoms can be severe and protracted.

How to relieve withdrawal symptoms

Returning to your previous dose will often relieve symptoms. When ready to try again, decrease your dose even more slowly, making smaller reductions and leaving longer between reductions, for example reducing dose by 2.5-5% every few weeks or months.

How to minimise withdrawal symptoms

The best way to minimise withdrawal symptoms is by slowly decreasing your dose, through to very low doses. This can take many months. You may think that the low doses in hyperbolic tapering are not needed, but small dose changes can have large effects. Do NOT skip the second half of the tapering plan. Do NOT skip a day or alternate days or doses, and do not suddenly stop.

Build a support team around you, as you may not recognise withdrawal symptoms in yourself. Look after yourself. Try physical activity and relaxation, sleep and eat well, limit alcohol and other drugs.

Why stop antidepressants?

For the health and quality of life benefits. You may feel more alert and energetic and you may begin to feel your emotions again. You may find it easier to manage your weight, and your sexual functioning may improve. You may prefer living your life free of antidepressant medication.

There are benefits in reducing your dose, even if you cannot stop. It is not a failure if you do not manage to stop. You can always try again at some point in the future, tapering even more slowly.

THE ROYAL NEW ZEALAND COLLEGE OF GENERAL PRACTITIONERS ENDORSED ACTIVITY CUM SCIENTIA CARITAS VALID UNTIL 31 JANUARY 2027 TE WHARE TOHU RATA O AOTEAROA

RACGP Accepted clinical resource

Research Lead: Professor Katharine Wallis. Further information is available via the RELEASE website: medical-school.uq.edu.au/release

Ethical approval: The University of Queensland Human Research Ethics Committee (2022/HE001667).

Funding: Commonwealth Department of Health, Medical Research Future Fund, Clinician Researchers Applied Research in Health - MRFAR000079. National Health and Medical Research Council (NHMRC) 2021 Partnership Projects PRC3 - 2015744.

© Copyright 2025 The University of Queensland. These materials are subject to copyright and may only be reproduced with permission of the University of Queensland. CRICOS Provider 00025B | TEQSA PRV12080. Reasonable care is taken to provide accurate information at the time of creation. This information is not intended as a substitute for medical advice. Read here for the Terms and Conditions.